## Direct observation of the Coulomb explosion in exotic atoms

<u>G.L. Borchert</u><sup>1</sup>, D. Anagnostopoulos<sup>2</sup>, D. Chatellard<sup>3</sup>, J.P. Egger<sup>3</sup>, P. El-Khoury<sup>4</sup>, D. Gotta<sup>1</sup>, P. Hauser<sup>5</sup>, P. Indelicato<sup>4</sup>, K. Kirch<sup>5</sup>, Th. Siems<sup>1</sup>, L.M. Simons<sup>5</sup>

<sup>1</sup> Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany
<sup>2</sup>Department of Material Science, University of Ioannina, GR-45110 Ioannina, Greece
<sup>3</sup>Institut de Physique de l'Université Neuchâtel, CH-2000 Neuchâtel, Switzerland
<sup>4</sup>Laboratoire Kastler-Brossel, UA 018 of the CNRS, Case 74,
Université Pierre et Marie Curie, 4 place Jussieu, F-75252 Paris Cedex 05, France
<sup>5</sup>Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland

In a series of experiments for a precision determination of the pion mass pionic and muonic X-ray transitions have been systematically investigated. The pions, produced in the accelerator facility of the PSI, Villigen, have been slowed down in the gas cell of a cyclotron trap to be captured by the gas atoms in highly excited bound atomic states. During the consequent radiative deexcitation processes characteristic X-rays were emitted, which are analyzed by a high resolution doubly focussing crystal spectrometer and registered in a optimized CCD detector system with high spatial granularity. Part of the pions decay to muons which undergo the analogue processes and are measured simultaneously. When comparing the spectra of di-atomic systems (Nitrogen,Oxygen) with those of mono-atomic systems (Ne) for the first time a broadening of the X-ray lines in the former case has been observed, which is attributed to the Doppler shift of the recoiling di-atomic Systems due to Coulomb explosion of the corresponding molecules. A study of the intensity dependence of parallel transitions as a function of pressure supports this interpretation. With a simple model the relevant charge states of the recoiling fragments have been estimated to be predominantly 3 or 4.