Dynamics of two interacting Bose condensates in a magnetostatic trap

M. Modugno, F. Dalfovo¹, P. Maddaloni², C. Fort, F. Minardi, and M. Inguscio

INFM - European Laboratory for Non Linear Spectroscopy, Università di Firenze,

Largo E. Fermi 2, I-50125 Firenze, Italy

Tel +39-055-2307809, Fax +39-055-224072

E-mail: modugno@fi.infn.it,

Dipartimento di Matematica e Fisica, Università Cattolica

Via Trieste 17, I-25121 Brescia, Italy

² Dipartimento di Fisica, Università di Padova, Via F. Marzolo 8, I-35131 Padova, Italy

We have induced collective oscillations of a ^{87}Rb $|F=2,m_F=2\rangle$ Bose condensate in a Ioffe-Pritchard type trap, by suddenly changing its number of atoms, hence its internal mean-field energy. A rf-pulse transfers 13 % of the atoms in the $|2,1\rangle$ state $(2 \cdot 10^4 \text{ out of } 1.5 \cdot 10^5)$ and a negligible population in the remaining sublevels. Due to the presence of gravity the equilibrium position of $|2,1\rangle$ is placed 9 μ m below that of $|2,2\rangle$ condensate. Once generated by the rf pulse, the $|2,1\rangle$ starts oscillating and undergoes periodical collisions with the $|2,2\rangle$. The latter represents, in a first approximation, just a deformation of the harmonic external potential experienced by the $|2,1\rangle$, thus causing a shift of its radial oscillation frequency ω_{1r} .

The main result of our work is a 5 % accurate measure of this frequency shift, that is as large as $0.06\omega_{1r}$. We have also measured the damping of this center-of-mass oscillation and the collective excitations of the $|2,2\rangle$ condensate via the observation of its aspect ratio as a function of its permanence time in the trap after the rf pulse.

These result have been compared with the predictions of a 2-dimensional model based on two coupled Gross-Pitaevskij (GP) equations, which captures the basic features of our system.