# Nuclear effects and neutron structure in deeply virtual Compton scattering off <sup>3</sup>He

Matteo Rinaldi & Sergio Scopetta

Dipartimento di Fisica dell'Università di Perugia and INFN, Sezione di Perugia, Italy.



Ottobre, 30 2013

## Outline

- Deeply virtual Compton scattering (DVCS) and nuclear Generalized Parton Distributions (GPDs)
  - GPDs of <sup>3</sup>He:
    - \* GPDs H, E, H in Impulse Approximation (IA) (S.Scopetta PRC 70 (2004) 015205; PRC 79 (2009) 025207)
    - \* Extracting the neutron information from <sup>3</sup>He data (M. R, S.Scopetta, PRC 85, 062201(R) (2012); PRC 87, 035208 (2013))
- Preliminary results for the GPD  $\tilde{H}$  of <sup>3</sup>He
- Conclusions



#### **GPDs - why?**

Initially, the interest in GPDs was a consequence of the "Spin crisis" (EMC, '88): most of the proton spin NOT carried by the quark helicities  $\Sigma$ 

Spin Sum Rule:

$$\sum + L_q + J_g = \frac{1}{2}$$

**OAM**  $(L_q)$  accessed through non forward processes:



DVCS -> GPDs



#### SiDIS -> TMDs



## GPDS: Definition (X. Ji PRL 78 (97) 610)



$$\int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) \quad \gamma^{\mu} \quad \psi_q(\lambda n/2) | P \rangle = H_q(x,\xi,\Delta^2) \bar{U}(P') \gamma^{\mu} U(P)$$
$$+ \quad E_q(x,\xi,\Delta^2) \bar{U}(P') \frac{i\sigma^{\mu\nu} \Delta_{\nu}}{2M} U(P) + \dots$$



## GPDS: Definition (X. Ji PRL 78 (97) 610)



 $\begin{aligned} \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) & \gamma^{\mu} \gamma_5 & \psi_q(\lambda n/2) | P \rangle = \tilde{H}_q(x,\xi,\Delta^2) \bar{U}(P') \gamma^{\mu} U(P) \\ &+ & \tilde{E}_q(x,\xi,\Delta^2) \bar{U}(P') \frac{\gamma_5 \Delta^{\mu}}{2M} U(P) + \dots \end{aligned}$ 



#### **GPDs: limits**

when P' = P, i.e.,  $\Delta^2 = \xi = 0$ , one recovers the usual PDFs:



 $H_q(x,0,0) = q(x);$   $\tilde{H}_q(x,0,0) = \Delta q(x);$   $E_q(x,0,0), \tilde{E}_q(x,0,0)$  unknown

the x-integration yields the q-contribution to the Form Factors (ffs)

$$\begin{split} \int dx \, \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) \gamma^\mu \psi_q(\lambda n/2) | P \rangle = \\ \int dx \, H_q(x,\xi,\Delta^2) \bar{U}(P') \gamma^\mu U(P) + \int dx \, E_q(x,\xi,\Delta^2) \bar{U}(P') \frac{\sigma^{\mu\nu} \Delta_\nu}{2M} U(P) + \dots \\ \Longrightarrow \int dx \, H_q(x,\xi,\Delta^2) = F_1^q(\Delta^2) \qquad \int dx \, E_q(x,\xi,\Delta^2) = F_2^q(\Delta^2) \end{split}$$

 $\tilde{G}_M^q = H_q + E_q$  one has  $\int dx \, \tilde{G}_M^q(x,\xi,\Delta^2) = G_M^q(\Delta^2)$ 



 $\implies$  Defining

#### **GPDs: limits**

when P' = P, i.e.,  $\Delta^2 = \xi = 0$ , one recovers the usual PDFs:



 $H_q(x,0,0) = q(x); \quad \tilde{H}_q(x,0,0) = \Delta q(x); \quad E_q(x,0,0), \tilde{E}_q(x,0,0) \text{ unknown}$ 

the x-integration yields the q-contribution to the Form Factors (ffs)

$$\begin{split} \int dx \, \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P' | \bar{\psi}_q(-\lambda n/2) \gamma^\mu \gamma_5 \psi_q(\lambda n/2) | P \rangle = \\ \int dx \, \tilde{H}_q(x,\xi,\Delta^2) \bar{U}(P') \gamma^\mu U(P) + \int dx \, \tilde{E}_q(x,\xi,\Delta^2) \bar{U}(P') \frac{\gamma_5 \Delta^\mu}{2M} U(P) + .. \\ \Longrightarrow \int dx \, \tilde{H}_q(x,\xi,\Delta^2) = g_A^q(\Delta^2) \qquad \int dx \, \tilde{E}_q(x,\xi,\Delta^2) = g_P^q(\Delta^2) \end{split}$$



#### **GPDs: A unique tool...**

- to explore the 3-dimensional structure of hadrons at parton level and for many other aspects...
- ...the most important here: access to the parton orbital angular momentum (OAM), solution (?) of the "Spin Crisis": **Ji Sum Rule**:

$$\langle J_q \rangle = \langle \Sigma_q \rangle + \langle L_q \rangle = \int_{-1}^1 dx \, x \, \tilde{G}_M^q(x, 0, 0)$$

#### ... but also an experimental challenge:

- Hard exclusive processes  $\longrightarrow$  small X-sections;
- Difficult extraction:

$$T_{\mathbf{DVCS}} \propto \int_{-1}^{1} dx \, \frac{H_q(x,\xi,\Delta^2)}{x-\xi+i\epsilon} + \dots \qquad ,$$



Competing **BH** process! Interference ( $\sigma$ -differences) measured.

$$d\sigma \propto |T_{\mathbf{DVCS}}|^2 + |T_{\mathbf{BH}}|^2 + 2 \Re\{T_{\mathbf{DVCS}}T^*_{\mathbf{BH}}\}$$

DVCS

 $\wedge \sim \sim$ 

BH

#### **Extracting GPDs**

One measures asymmetries:  $A = \frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-}$ 



Polarized beam, unpolarized target:

$$\Delta \sigma_{LU} \simeq \sin \phi \left[ F_1 \mathbf{H} + \xi (F_1 + F_2) \tilde{\mathbf{H}} + (\Delta^2 F_2 / M^2) \mathbf{E} / 4 \right] d\phi \quad \Longrightarrow \quad \mathbf{H}$$

Unpolarized beam, longitudinally polarized target:  $\Delta \sigma_{UL} \simeq \sin \phi \left\{ F_1 \tilde{\mathbf{H}} + \xi (F_1 + F_2) \left[ \mathbf{H} + \xi / (1 + \xi) \mathbf{E} \right] \right\} d\phi \qquad \Longrightarrow \qquad \tilde{\mathbf{H}}$ 

Unpolarized beam, transversely polarized target:

$$\Delta \sigma_{UT} \simeq \cos \phi \sin(\phi_S - \phi) \left[ \Delta^2 (F_2 \mathbf{H} - F_1 \mathbf{E}) / M^2 \right] d\phi \implies \mathbf{E}$$



To evaluate cross sections, e.g. for experiments planning, one needs H, H, EThis is what we have calculated for <sup>3</sup>He. Why nuclei? Why <sup>3</sup>He?

Ottobre, 30 2013

#### Why nuclei?

#### Several relevant issues can be investigated...:

- the nuclear short range structure, at quark level, can be accessed and the reaction mechanism of DIS off nuclei, e.g. the validity of I.A. and the relevance of effects beyond it (non nucleonic degrees of freedom, nucleon modifications...) can be investigated... origin of the EMC effect...
- very important here: the neutron, always from nuclear targets
- ... with some effort: measurements are difficult:
  - In principle, need for a recoil detector to be sure that the nucleus did not break (despite of this, some data are already available!);
  - Few data already available:

Airapetian et al. (Hermes) NPB 829, 1 (2010); PRC 81, 035202 (2010)  $D(\vec{e}, e'\gamma)X$ ,  $\vec{D}(\vec{e}, e'\gamma)X$ ,  $Ne(\vec{e}, e'\gamma)X$ ; and then He, N, Ne, Kr, Xe... Little *A*-dependence found



Mazouz et al. (JLab Hall A) PRL 99.242501 (2007); experiment E08-025 (2010)  $D(\vec{e}, e'\gamma)X = d(\vec{e}, e'\gamma)d + n(\vec{e}, e'\gamma)X + p(\vec{e}, e'\gamma)X$ ; <sup>3</sup>He target?

#### GPDs for <sup>3</sup>He: why?

- <sup>3</sup>He is theoretically well known. Even a relativistic treatment may be implemented.
- <sup>3</sup>He has been used extensively as an effective neutron target, especially to unveil the spin content of the free neutron, due to its peculiar spin structure:



<sup>3</sup>He always promising when the neutron angular momentum properties have to be studied. To what extent for OAM and  $\tilde{G}_M^q$ ? The answer here.

#### To this aim, <sup>3</sup>He is a unique target:

\*

in isoscalar systems, such as <sup>2</sup>H and <sup>4</sup>He, the contribution of the neutron  $E_q$  is basically cancelled by that of the proton one ( $\kappa_p \simeq -\kappa_n$ ); vey difficult to extract the neutron  $E_q$ , crucial to access OAM, in coherent experiments;

\*

heavier targets do not allow refined theoretical treatments.



coherent DVCS in I.A. (<sup>3</sup>He does not break-up,  $\Delta^2 \ll M^2, \xi^2 \ll 1$ , ):

In a symmetric frame (  $\bar{p}=(p+p^\prime)/2$  ) :

$$k^{+} = (x+\xi)\bar{P}^{+} = (x'+\xi')\bar{p}^{+} ,$$
  
$$(x+\Delta)^{+} = (x-\xi)\bar{P}^{+} = (x'-\xi')\bar{p}^{+} ,$$

one has, for a given GPD,  $H_q$ ,  $\tilde{H}_q$  or  $\tilde{G}^q_M$ ,

$$GPD_{q}(x,\xi,\Delta^{2}) \simeq \int \frac{dz^{-}}{4\pi} e^{ix\bar{P}^{+}z^{-}}{}_{A} \langle P'S' | \hat{O}_{q}^{\mu} | PS \rangle_{A} |_{z^{+}=0, z_{\perp}=0} \,.$$



coherent DVCS in I.A. (<sup>3</sup>He does not break-up,  $\Delta^2 \ll M^2, \xi^2 \ll 1$ , ):

In a symmetric frame (  $\bar{p}=(p+p')/2$  ) :

$$k^{+} = (x+\xi)\bar{P}^{+} = (x'+\xi')\bar{p}^{+} ,$$
  
$$(x+\Delta)^{+} = (x-\xi)\bar{P}^{+} = (x'-\xi')\bar{p}^{+} ,$$

one has, for a given GPD,  $H_q$ ,  $\tilde{H}_q$  or  $\tilde{G}^q_M$ ,

$$GPD_q(x,\xi,\Delta^2) \simeq \int \frac{dz^-}{4\pi} e^{ix\bar{P}^+z^-} {}_A \langle P'S' | \hat{O}^{\mu}_q | PS \rangle_A |_{z^+=0, z_\perp=0} \cdot$$

By properly inserting complete sets of states for the interacting nucleon and the recoiling system :



coherent DVCS in I.A. (<sup>3</sup>He does not break-up,  $\Delta^2 \ll M^2, \xi^2 \ll 1$ , ):

 $\begin{array}{c} \mathbf{e} \\ \gamma^{*} \mathbf{q} \\ \mathbf{p} \\ \mathbf{p$ 

e

In a symmetric frame (  $\bar{p}=(p+p^\prime)/2$  ) :

$$k^{+} = (x+\xi)\bar{P}^{+} = (x'+\xi')\bar{p}^{+} ,$$
  
$$k+\Delta)^{+} = (x-\xi)\bar{P}^{+} = (x'-\xi')\bar{p}^{+} ,$$

one has, for a given GPD, 
$$H_q$$
,  $\tilde{H}_q$  or  $\tilde{G}_M^q$ ,  

$$GPD_q(x,\xi,\Delta^2) = \int \frac{dz^-}{4\pi} e^{ix'\bar{p}^+z^-} \langle P'S'| \sum_{\vec{P}'_R,S'_R,\vec{p}',s'} \{|P'_RS'_R\rangle|p's'\rangle\} \langle P'_RS'_R|$$

$$\langle p's'|\hat{O}_q^{\mu} \sum_{\vec{P}_R,S_R,\vec{p},s} \{|P_RS_R\rangle|ps\rangle\} \{\langle P_RS_R|\langle ps|\} |PS\rangle,$$

and, since 
$$\{\langle P_R S_R | \langle ps | \} | PS \rangle = \langle P_R S_R, ps | PS \rangle (2\pi)^3 \delta^3 (\vec{P} - \vec{P}_R - \vec{p}) \delta_{S,S_R s}$$
,



 $H_q^A$  can be obtained in terms of  $H_q^N$  (S.Scopetta PRC 70, 015205 (2004), PRC 79, 025207 (2009)):

$$H_q^A(x,\xi,\Delta^2) = \sum_N \int dE \int d\vec{p} \,\overline{\sum_S} \sum_s P_{SS,ss}^N(\vec{p},\vec{p}',E) \frac{\xi'}{\xi} H_q^N(x',\Delta^2,\xi') ,$$

 $\tilde{G}_{M}^{3,q}$  in terms of  $\tilde{G}_{M}^{N,q}$  (M. R, S.Scopetta PRC 85, 062201(R) (2012); PRC 87, 035208 (2013)):

$$\tilde{G}_{M}^{3,q}(x,\Delta^{2},\xi) = \sum_{N} \int dE \int d\vec{p} \left[ P_{+-,+-}^{N} - P_{+-,-+}^{N} \right] (\vec{p},\vec{p}',E) \frac{\xi'}{\xi} \tilde{G}_{M}^{N,q}(x',\Delta^{2},\xi') ,$$

and  $\tilde{H}_q^A$  can be obtained in terms of  $\tilde{H}_q^N$  (preliminary):

$$\tilde{H}_{q}^{A}(x,\xi,\Delta^{2}) = \sum_{N} \int dE \int d\vec{p} \left[ P_{++,++}^{N} - P_{++,--}^{N} \right] (\vec{p},\vec{p}',E) \frac{\xi'}{\xi} \tilde{H}_{q}^{N}(x',\Delta^{2},\xi') ,$$



 $H_q^A$  can be obtained in terms of  $H_q^N$  (S.Scopetta PRC 70, 015205 (2004), PRC 79, 025207 (2009)):

$$H_q^A(x,\xi,\Delta^2) = \sum_N \int dE \int d\vec{p} \,\overline{\sum_S} \sum_s P_{SS,ss}^N(\vec{p},\vec{p}',E) \frac{\xi'}{\xi} H_q^N(x',\Delta^2,\xi') ,$$

 $\tilde{G}_{M}^{3,q}$  in terms of  $\tilde{G}_{M}^{N,q}$  (M. R, S.Scopetta PRC 85, 062201(R) (2012); PRC 87, 035208 (2013)):

$$\tilde{G}_{M}^{3,q}(x,\Delta^{2},\xi) = \sum_{N} \int dE \int d\vec{p} \left[ P_{+-,+-}^{N} - P_{+-,-+}^{N} \right] (\vec{p},\vec{p}',E) \frac{\xi'}{\xi} \tilde{G}_{M}^{N,q}(x',\Delta^{2},\xi') ,$$

where  $P_{SS,ss}^{N}(\vec{p}, \vec{p'}, E)$  is the one-body, spin-dependent, off-diagonal spectral function for the nucleon N in the nucleus,

$$P^{N}_{SS',ss'}(\vec{p},\vec{p}',E) = \frac{1}{(2\pi)^6} \frac{M\sqrt{ME}}{2} \int d\Omega_t \sum_{s_t} \langle \vec{P'}S' | \vec{p}'s', \vec{t}s_t \rangle_N \langle \vec{p}s, \vec{t}s_t | \vec{P}S \rangle_N ,$$

evaluated by means of a realistic treatment based on Av18 wave functions (w.f. from A. Kievsky *et al* NPA 577, 511 (1994), overlaps from A. Kievsky *et. al*, PRC 56, 64 (1997)).

Nucleon GPDs: initially, a simple model by Radyushkin&Musatov (PRD 61, 074027 (2000)) Nuclear effects and neutron structure in deeply virtual Compton scattering off <sup>3</sup> He - p.11/23

## The $\tilde{G}_M^{3,q}$ calculation has the correct limits:

For  $H_q^3$ , correct forward limit and *x*-integral (S.Scopetta PRC 70, (2004), PRC 79, (2009)); For  $\tilde{G}_M^3$  (M. R, S.Scopetta PRC 85, 062201(R) (2012); PRC 87, 035208 (2013) ):

1 - Forward limit: no control on  $E_q^3(x,0,0)$  no possible check;

2 - Magnetic F.F.:

 $\sum_q \int dx \, \tilde{G}^{3,q}_M(x,\xi,\Delta^2) = G^3_M(\Delta^2)$ 

- in perfect agreement with previous IA, Av18 calculations (L.E. Marcucci et al. PRC 58 (1998))
  - in good agreement with data in the region relevant to the coherent process,  $-\Delta^2 \ll 0.15 \text{ GeV}^2$
- To have agreement at higher  $\Delta^2$ , effects beyond IA are necessary: not important for the coherent channel!



### $\tilde{G}_M^{3,q}$ : proton and neutron contributions

1 - Forward limit,  $\Delta^2 = 0, \xi = 0$ :

As we hoped, the neutron contribution to <sup>3</sup>He largely dominates!  $(x_3 = (M_A/M)x \simeq 3x)$ : The proton contribution to <sup>3</sup>He is almost negligible!

2 - Non-forward,  $\Delta^2=-0.1~{\rm GeV^2}, \xi=0.1$ :

The neutron contribution to <sup>3</sup>He still dominates The proton contribution to <sup>3</sup>He gets sizable

How to get the neutron information?



## $\tilde{G}_M^{3,q}$ : Flavor separation

For the u flavor, the neutron contribution (dashed) to <sup>3</sup>He (full) is less important than for the d flavor:



Understandable, sketching the formula:

 $\tilde{G}_M^{3,q} \approx P_p^3 \otimes \tilde{G}_M^{p,q} + P_n^3 \otimes \tilde{G}_M^{n,q} ,$ 

where  $P_{p(n)}^3$  describes the proton (neutron) dynamics in <sup>3</sup>He.

As already explained, due to the spin structure of <sup>3</sup>He,  $P_n^3 >> P_p^3 \longrightarrow$  neutron dominates in the forward limit.

With increasing  $\Delta^2$ , for the  $\mathcal{U}$  flavor,  $\tilde{G}_M^{p,u} >> \tilde{G}_M^{n,u} \longrightarrow$  the proton contribution grows. Not for d!

Besides, 1/2 of the *d* content of <sup>3</sup>He comes from the neutron, only 1/5 of the u one comes from it.



#### **Extracting the neutron - I:**

The convolution formula can be written as

$$\tilde{G}_{M}^{3,q}(x_{3},\Delta^{2},\xi) = \sum_{N} \int_{x_{3}}^{\frac{M_{A}}{M}} \frac{dz}{z} g_{N}^{3}(z,\Delta^{2},\xi) \tilde{G}_{M}^{N,q}\left(\frac{x_{3}}{z},\Delta^{2},\frac{\xi}{z},\right) ,$$

where  $g_N^3(z, \Delta^2, \xi)$  is a "light cone off-forward momentum distribution" and, since close to the forward limit it is strongly peaked around z = 1

$$g_N^3(z,\Delta^2,\xi) = \int dE \int d\vec{p} \, \tilde{P}_N^3(\vec{p},\vec{p}+\vec{\Delta},E)$$
  $\delta\left(z+\xi-rac{M_A}{M}rac{p^+}{ar{p}^+}
ight)$ 





Ottobre, 30 2013

#### **Extracting the neutron - I:**

The convolution formula can be written as

$$\tilde{G}_{M}^{3,q}(x_{3},\Delta^{2},\xi) = \sum_{N} \int_{x_{3}}^{\frac{M_{A}}{M}} \frac{dz}{z} g_{N}^{3}(z,\Delta^{2},\xi) \tilde{G}_{M}^{N,q}\left(\frac{x_{3}}{z},\Delta^{2},\frac{\xi}{z},\right) ,$$

where  $g_N^3(z, \Delta^2, \xi)$  is a "light cone off-forward momentum distribution" and, since close to the forward limit it is strongly peaked around z = 1

$$\begin{split} \tilde{G}_M^{3,q}(x_3,\Delta^2,\xi) &\simeq low\,\Delta^2 \simeq \sum_N \tilde{G}_M^{N,q}\left(x_3,\Delta^2,\xi\right) \int_0^{\frac{M_A}{M}} dz g_N^3(z,\Delta^2,\xi) \\ &= G_M^{3,p,point}(\Delta^2) \tilde{G}_M^p(x_3,\Delta^2,\xi) + G_M^{3,n,point}(\Delta^2) \tilde{G}_M^n(x_3,\Delta^2,\xi) \,. \end{split}$$

where, at  $x_3 < 0.7$ , the magnetic point like ff has been introduced

$$G_{M}^{3,N,point}(\Delta^{2}) = \int dE \int d\vec{p} \, \vec{P}_{N}^{3}(\vec{p},\vec{p}+\vec{\Delta},E) = \int_{0}^{\frac{M_{A}}{M}} dz \, g_{N}^{3}(z,\Delta^{2},\xi) \; .$$



#### Extracting the neutron - II:

Validity of the approximated formula: full: IA calculation,  $\tilde{G}^3_M(x,\Delta^2,\xi)$  and proton and neutron contributions to it, at  $\Delta^2 = -0.1 \text{ GeV}^2$ ,  $\xi = 0.1$ ;

dashed: same quantities, with the approximated formula:



Impressive agreement! The only Nuclear Physics ingredient in the approximated formula is the magnetic point like ff, which is under good theoretical control:

$\Delta^2$	$G_M^{3,p,point}$	$G_M^{3,p,point}$	$G_M^{3,n,point}$	$G_M^{3,n,point}$
[GeV <sup>2</sup> ]	Av18	Av14	Av18	Av14
0	-0.044	-0.049	0.879	0.874
-0.1	0.040	0.038	0.305	0.297
-0.2	0.036	0.035	0.125	0.119



#### **Extracting the neutron - III:**

The approximated relation can now be solved to extract the neutron contribution:

$$\begin{split} \tilde{G}_M^{n,extr}(x,\Delta^2,\xi) &\simeq \quad \frac{1}{G_M^{3,n,point}(\Delta^2)} \left\{ \tilde{G}_M^3(x,\Delta^2,\xi) \right. \\ &- \quad G_M^{3,p,point}(\Delta^2) \tilde{G}_M^p(x,\Delta^2,\xi) \right\} \,, \end{split}$$

from data for  $\tilde{G}^3_M(x, \Delta^2, \xi)$  and  $\tilde{G}^p_M(x, \Delta^2, \xi)$ , using as theoretical ingredients the magnetic point like ffs only.

The procedure works nicely!

full : the neutron model for  $\tilde{G}_M^n(x, \Delta^2, \xi)$ and the different flavor contributions to it used in the IA calculation, at  $\Delta^2 = -0.1 \text{ GeV}^2$ ,  $\xi = 0.1$ ;

dashed: the neutron extracted using the IA calculation for  $\tilde{G}_M^3(x, \Delta^2, \xi)$ and the model used in it for  $\tilde{G}_M^p(x, \Delta^2, \xi)$ together with the magnetic point like ffs.





#### **Extracting the neutron - IV:**

The validity of the extraction procedure is emphasized showing the following ratio, which would be one if the procedure were perfect:



at x < 0.7, in all the kinematical range relevant for coherent DVCS at JLab, the error in the extraction is a few percents.



#### **Extracting the neutron - V:**

The validity of the extraction procedure is emphasized showing the same ratio, evaluated using different models for the GPDs as input in the IA calculation:

$$r_n(x,\Delta^2,\xi) = \frac{\tilde{G}_M^{n,extr}(x,\Delta^2,\xi)}{\tilde{G}_M^n(x,\Delta^2,\xi)}$$

dashed: the model of Radyushkin, at  $\Delta^2 = -0.1 \ {\rm GeV}^2, \ \xi = 0 \ ; \label{eq:Lagrangian}$ 

full: a very different model based on a constituent quark scenario (S.Scopetta, V. Vento EPJA 16, 527 (2003) ) at  $\Delta^2 = -0.1 \text{ GeV}^2, \xi = 0$ ;

crosses: another very different model, the MIT bag model (X.-D. Ji et al., PRD 56 (1997) 5511 ) at  $\Delta^2=-0.1~{\rm GeV}^2,\,\xi=0~.$ 





at x < 0.7, in all the kinematical range relevant for coherent DVCS at JLab, the error in the extraction due to the use of a different nucleonic model is a few percents at most.

#### **The GPD** $\tilde{H}$ : preliminary

 $\tilde{H}^{3,u}(x,\Delta^2,\xi)$  and proton and (dominant!) neutron contributions to it:



full: IA calculation; dashed: approximated formula:

$$\tilde{H}^{3,u}(x,\Delta^2,\xi) \simeq g_A^{3,p,point}(\Delta^2)\tilde{H}^{p,u}(x,\Delta^2,\xi) + g_A^{3,n,point}(\Delta^2)\tilde{H}^{n,u}(x,\Delta^2,\xi)$$

Good agreement! The only Nuclear Physics ingredient in the approximated formula is the axial point like ff, which is under good theoretical control. One has  $g_A^{3,N,point}(\Delta^2 = 0) = p_N$ , nucleon effective polarizations (within AV18,  $p_n = 0.878, p_p = -0.024$ ), used in DIS for extracting the neutron information from <sup>3</sup>He (C. Ciofi, S.Scopetta, E. Pace and G. Salmè, PRC 48 R968 (1993)).



## Conclusions

What we have:

- \* An instant form, I.A. calculation of  $H^3$ ,  $\tilde{G}^3_M$ ,  $\tilde{H}^3$ , within AV18;
- \* the neutron contribution dominates  $\tilde{G}_M^3$  and  $\tilde{H}^3$  at low  $\Delta^2$ ;
- an extraction procedure of the neutron information, able to take into account all the nuclear effects encoded in an IA analysis, weakly dependent on both the nuclear potential and the nucleonic model used in the calculation;
- What we can do now: to estimate X-sections (DVCS, BH, Interference)  $\rightarrow$  a proposal of coherent DVCS off <sup>3</sup>He at JLab@12 GeV?
- What has to be done, in case experiments are performed at higher  $\Delta^2$ :
  - \* To implement a **RELATIVISTIC TREATMENT**
  - f and/or to go beyond IA, including many body currents into the scheme.

#### What is expected:

- \* The OAM parton structure of the neutron;
- \* A deeper understanding of the nuclear parton structure;

#### Importance of nuclear FB systems for QCD studies



#### coherent vs. incoherent DVCS:





Coherent





Incoherent



Ottobre,  $30\,2013$ 

#### A few words about $P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E)$ :

$$P_N^3(\vec{p}, \vec{p} + \vec{\Delta}, E) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_M \sum_{f,s} \langle \vec{P}' M | (\vec{P} - \vec{p}) S_f, (\vec{p} + \vec{\Delta}) s \rangle$$
  
 
$$\times \quad \langle (\vec{P} - \vec{p}) S_f, \vec{ps} | \vec{P} M \rangle \, \delta(E - E_{min} - E_f^*) \, .$$



- the two-body recoiling system can be either the deuteron or a scattering state;
- when a deeply bound nucleon, with high removal energy  $E = E_{min} + E_f^*$ , leaves the nucleus, the recoling system is left with high excitation energy  $E_f^*$ ;
- the three-body bound state and the two-body bound or scattering state are evaluated within the same (Av18) interaction: the extension of the treatment to heavier nuclei would be extremely difficult